

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use tranexamic acid USP tablets safely and effectively. See full prescribing information for tranexamic acid USP tablets.

Tranexamic Acid, USP Tablets
Initial U.S. Approval: 1986

INDICATIONS AND USAGE

Tranexamic Acid USP Tablets is an antifibrinolytic indicated for the treatment of cyclic heavy menstrual bleeding.(1)

DOSAGE AND ADMINISTRATION

- 1,300 mg (two 650 mg tablets) three times a day (3,900 mg/day) for a maximum of 5 days during monthly menstruation (2.1)
- Renal impairment: Dosage adjustment is needed if serum creatinine concentration (Cr) is higher than 1.4 mg/dL (2.2)
 - Cr above 1.4 mg/dL and ≤ 2.8 mg/dL: 1,300 mg (two 650 mg tablets) two times a day (2,600 mg/day) for a maximum of 5 days during menstruation
 - Cr above 2.8 mg/dL and ≤ 5.7 mg/dL: 1,300 mg (two 650 mg tablets) once a day (1,300 mg/day) for a maximum of 5 days during menstruation
 - Cr above 5.7 mg/dL: 650 mg (one 650 mg tablet) once a day (650 mg/day) for a maximum of 5 days during menstruation

DOSAGE FORMS AND STRENGTHS

Tablets: 650 mg (3)

CONTRAINDICATIONS

- Women who are using combination hormonal contraception (4.1)
- Women with active thromboembolic disease or a history or intrinsic risk of thrombosis or thromboembolism, including retinal vein or artery occlusion (4.1)
- Hypersensitivity to tranexamic acid (4.2)

WARNINGS AND PRECAUTIONS

- Concomitant use of tranexamic acid USP tablets with Factor IX complex concentrates, anti-inhibitor coagulant concentrates or all-trans retinoic acid (oral tretinoin) may increase the risk of thrombosis. (5.1)
- Visual or ocular adverse effects may occur with tranexamic acid USP tablets. Immediately discontinue use if visual or ocular symptoms occur. (5.1)
 - In case of severe allergic reaction, discontinue tranexamic acid USP tablets and seek immediate medical attention. (5.2)
 - Cerebral edema and cerebral infarction may be caused by use of tranexamic acid USP tablets in women with subarachnoid hemorrhage. (5.3)
 - Ligneous conjunctivitis has been reported in patients taking tranexamic acid. (5.4)

ADVERSE REACTIONS

Most common adverse reactions in clinical trials (≥ 5%, and more frequent in tranexamic acid USP tablets subjects compared to placebo subjects) are headache, sinus and nasal symptoms, back pain, abdominal pain, musculoskeletal pain, joint pain, muscle cramps, migraine, anemia and fatigue. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Amring Pharmaceuticals Inc. at 1-844-Amring1 (1-844-267-4641) or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

Concomitant therapy with tissue plasminogen activators may decrease the efficacy of both tranexamic acid USP tablets and tissue plasminogen activators. (7.2)

USE IN SPECIFIC POPULATIONS

- Geriatric Use: Tranexamic Acid USP Tablets is not indicated for use in postmenopausal women (8.5)
- Renal impairment: Dosage adjustment is needed. (2.2, 8.6)
- Hepatic impairment: No dosage adjustment is needed. (8.7)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Rev. 03/2019

FULL PRESCRIBING INFORMATION: CONTENTS*

- INDICATIONS AND USAGE
 - DOSAGE AND ADMINISTRATION
 - Recommended Dosage
 - Renal Impairment
 - DOSAGE FORMS AND STRENGTHS
 - CONTRAINDICATIONS
 - Thromboembolic Risk
 - Hypersensitivity to Tranexamic Acid
 - WARNINGS AND PRECAUTIONS
 - Thromboembolic Risk
 - Severe Allergic Reaction
 - Subarachnoid Hemorrhage
 - Ligneous Conjunctivitis
 - ADVERSE REACTIONS
 - Clinical Trial Experience
 - Postmarketing Experience
 - DRUG INTERACTIONS
 - Hormonal Contraceptives
 - Tissue Plasminogen Activators
 - Factor IX Complex Concentrates or Anti-Inhibitor Coagulant Concentrates
 - All-Trans Retinoic Acid (Oral Tretinoin)
 - USE IN SPECIFIC POPULATIONS
 - Pregnancy
 - Nursing Mothers
 - Pediatric Use
 - Geriatric Use
 - Renal Impairment
 - Hepatic Impairment
 - OVERDOSAGE
 - DESCRIPTION
 - CLINICAL PHARMACOLOGY
 - Mechanism of Action
 - Pharmacodynamics
 - Pharmacokinetics
 - NONCLINICAL TOXICOLOGY
 - Carcinogenesis, Mutagenesis, Impairment of Fertility
 - Animal Toxicology and/or Pharmacology
 - CLINICAL STUDIES
 - Three-Cycle Treatment Study
 - Six-Cycle Treatment Study
 - MBL Results over Time
 - HOW SUPPLIED/STORAGE AND HANDLING
 - PATIENT COUNSELING INFORMATION
- *Sections or subsections omitted from the full prescribing information are not listed

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Tranexamic Acid USP Tablets are indicated for the treatment of cyclic heavy menstrual bleeding [see *Clinical Studies* (14)].

Prior to prescribing tranexamic acid USP tablets, exclude endometrial pathology that can be associated with heavy menstrual bleeding.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage

The recommended dose of tranexamic acid USP tablets for women with normal renal function is two 650 mg tablets taken three times daily (3900 mg/day) for a maximum of 5 days during monthly menstruation. Tranexamic Acid USP Tablets may be administered without regard to meals. Tablets should be swallowed whole and not chewed or broken apart.

2.2 Renal Impairment

In patients with renal impairment, the plasma concentration of tranexamic acid increased as serum creatinine concentration increased [see *Clinical Pharmacology* (12.3)]. Dosage adjustment is needed in patients with serum creatinine concentration higher than 1.4 mg/dL (Table 1).

Table 1. Dosage of tranexamic acid USP tablets in Patients with Renal Impairment

Tranexamic Acid USP Tablets		
Serum Creatinine (mg/dL)	Adjusted Dose	Total Daily Dose
Cr above 1.4 and ≤ 2.8	1300 mg (two 650 mg tablets) two times a day for a maximum of 5 days during menstruation	2600 mg
Cr above 2.8 and ≤ 5.7	1300 mg (two 650 mg tablets) once a day for a maximum of 5 days during menstruation	1300 mg
Cr above 5.7	650 mg (one 650 mg tablet) once a day for a maximum of 5 days during menstruation	650 mg

3 DOSAGE FORMS AND STRENGTHS

650 mg tablets

4 CONTRAINDICATIONS

4.1 Thromboembolic Risk

Do not prescribe tranexamic acid USP tablets to women who are

- using combination hormonal contraception
- known to have any of the following conditions:
 - Active thromboembolic disease (e.g., deep vein thrombosis, pulmonary embolism, or cerebral thrombosis)
 - A history of thrombosis or thromboembolism, including retinal vein or artery occlusion
 - An intrinsic risk of thrombosis or thromboembolism (e.g., thrombogenic valvular disease, thrombogenic cardiac rhythm disease, or hypercoagulopathy)

Venous and arterial thrombosis or thromboembolism, as well as cases of retinal artery and retinal vein occlusions, have been reported with tranexamic acid.

4.2 Hypersensitivity to Tranexamic Acid

Do not prescribe tranexamic acid USP tablets to women with known hypersensitivity to tranexamic acid [see *Warnings and Precautions* (5.2) and *Adverse Reactions* (6.1)].

5 WARNINGS AND PRECAUTIONS

5.1 Thromboembolic Risk

Concomitant Use of Hormonal Contraceptives

Combination hormonal contraceptives are known to increase the risk of venous thromboembolism, as well as arterial thromboses such as stroke and myocardial infarction. Because tranexamic acid USP tablets are antifibrinolytic, the risk of venous thromboembolism, as well as arterial thromboses such as stroke, may increase further when hormonal contraceptives are administered with tranexamic acid USP tablets. This is of particular concern in women who are obese or smoke cigarettes, especially smokers over 35 years of age.

Women using hormonal contraception were excluded from the clinical trials supporting the safety and efficacy of tranexamic acid USP tablets, and there are no clinical trial data on the risk of thrombotic events with the concomitant use of tranexamic acid USP tablets with hormonal contraceptives. However, there have been US postmarketing reports of venous and arterial thrombotic events in women who have used tranexamic acid USP tablets concomitantly with combination hormonal contraceptives. For this reason, concomitant use of tranexamic acid with combination hormonal contraceptives is contraindicated. [see *Contraindications* (4.1) and *Drug Interactions* (7.1)].

Factor IX Complex Concentrates or Anti-Inhibitor Coagulant Concentrates

Tranexamic Acid USP Tablets are not recommended for women taking either Factor IX complex concentrates or anti-inhibitor coagulant concentrates because the risk of thrombosis may be increased [see *Drug Interactions* (7.3) and *Clinical Pharmacology* (12.3)].

All-Trans Retinoic Acid (Oral Tretinoin)

Exercise caution when prescribing tranexamic acid USP tablets to women with acute promyelocytic leukemia taking all-trans retinoic acid for remission induction because of possible exacerbation of the procoagulant effect of all-trans retinoic acid [see *Drug Interactions* (7.4) and *Clinical Pharmacology* (12.3)].

Ocular Effects

Retinal venous and arterial occlusion has been reported in patients using tranexamic acid. Patients should be instructed to report visual and ocular symptoms promptly. In the event of such symptoms, patients should be instructed to discontinue tranexamic acid USP tablets immediately and should be referred to an ophthalmologist for a complete ophthalmic evaluation, including dilated retinal examination, to exclude the possibility of retinal venous or arterial occlusion.

5.2 Severe Allergic Reaction

A case of severe allergic reaction to tranexamic acid USP tablets was reported in the clinical trials, involving a subject who experienced dyspnea, tightening of her throat, and facial flushing that required emergency medical treatment. A case of anaphylactic shock has also been reported in the literature, involving a patient who received an intravenous bolus of tranexamic acid.

5.3 Subarachnoid Hemorrhage

Cerebral edema and cerebral infarction may be caused by use of tranexamic acid USP tablets in women with subarachnoid hemorrhage.

5.4 Ligneous Conjunctivitis

Ligneous conjunctivitis has been reported in patients taking tranexamic acid. The conjunctivitis resolved following cessation of the drug.

6 ADVERSE REACTIONS

6.1 Clinical Trial Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to the rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

Short-term Studies

The safety of tranexamic acid USP tablets in the treatment of heavy menstrual bleeding (HMB) was studied in two randomized, double-blind, placebo-controlled studies [see *Clinical Studies (14)*]. One study compared the effects of two doses of tranexamic acid USP tablets (1950 mg and 3900 mg given daily for up to 5 days during each menstrual period) versus placebo over a 3-cycle treatment duration. A total of 304 women were randomized to this study, with 115 receiving at least one dose of 3900 mg/day of tranexamic acid USP tablets. A second study compared the effects of tranexamic acid USP tablets (3900 mg/day) versus placebo over a 6-cycle treatment duration. A total of 196 women were randomized to this study, with 117 receiving at least one dose of tranexamic acid USP tablets. In both studies, subjects were generally healthy women who had menstrual blood loss of ≥ 80 mL.

In these studies, subjects were 18 to 49 years of age with a mean age of approximately 40 years, had cyclic menses every 21-35 days, and a BMI of approximately 32 kg/m². On average, subjects had a history of HMB for approximately 10 years and 40% had fibroids as determined by transvaginal ultrasound. Approximately 70% were Caucasian, 25% were Black, and 5% were Asian, Native American, Pacific Islander, or Other. Seven percent (7%) of all subjects were of Hispanic origin. Women using hormonal contraception were excluded from the trials.

The rates of discontinuation due to adverse events during the two clinical trials were comparable between tranexamic acid USP tablets and placebo. In the 3-cycle study, the rate in the 3900 mg tranexamic acid USP tablets dose group was 0.8% as compared to 1.4% in the placebo group. In the 6-cycle study, the rate in the tranexamic acid USP tablets group was 2.4% as compared to 4.1% in the placebo group. Across the studies, the combined exposure to 3900 mg/day tranexamic acid USP tablets was 947 cycles and the average duration of use was 3.4 days per cycle.

A list of adverse events occurring in $\geq 5\%$ of subjects and more frequently in tranexamic acid USP tablets treated subjects receiving 3900 mg/day compared to placebo is provided in Table 2.

Table 2. Adverse Events Reported by $\geq 5\%$ of Subjects Treated with tranexamic acid USP tablets and More Frequently in tranexamic acid USP tablets-treated Subjects

	Tranexamic Acid 3900 mg/day n (%) (N=232)	Placebo n (%) (N=139)
Total Number of Adverse Events	1500	923
Number of Subjects with at Least One Adverse Event	208 (89.7%)	122 (87.8%)
HEADACHE ^a	117 (50.4%)	65 (46.8%)
NASAL & SINUS SYMPTOMS ^b	59 (25.4%)	24 (17.3%)
BACK PAIN	48 (20.7%)	21 (15.1%)
ABDOMINAL PAIN ^c	46 (19.8%)	25 (18.0%)
MUSCULOSKELETAL PAIN ^d	26 (11.2%)	4 (2.9%)
ARTHRALGIA ^e	16 (6.9%)	7 (5.0%)
MUSCLE CRAMPS & SPASMS	15 (6.5%)	8 (5.8%)
MIGRAINE	14 (6.0%)	8 (5.8%)
ANEMIA	13 (5.6%)	5 (3.6%)
FATIGUE	12 (5.2%)	6 (4.3%)

^aIncludes headache and tension headache

^bNasal and sinus symptoms include nasal, respiratory tract and sinus congestion, sinusitis, acute sinusitis, sinus headache, allergic sinusitis and sinus pain, and multiple allergies and seasonal allergies

^cAbdominal pain includes abdominal tenderness and discomfort

^dMusculoskeletal pain includes musculoskeletal discomfort and myalgia

^eArthralgia includes joint stiffness and swelling

Long-term Studies

Long-term safety of tranexamic acid USP tablets was studied in two open-label studies. In one study, subjects with physician-diagnosed heavy menstrual bleeding (not using the alkaline hematin methodology) were treated with 3900 mg/day for up to 5 days during each menstrual period for up to 27 menstrual cycles. A total of 781 subjects were enrolled and 239 completed the study through 27 menstrual cycles. A total of 12.4% of the subjects withdrew due to adverse events. Women using hormonal contraception were excluded from the study. The total exposure in this study to 3900 mg/day tranexamic acid USP tablets was 10,213 cycles. The average duration of tranexamic acid USP tablets use was 2.9 days per cycle.

A long-term open-label extension study of subjects from the two short-term efficacy studies was also conducted in which subjects were treated with 3900 mg/day for up to 5 days during each menstrual period of up to 9 menstrual cycles. A total of 288 subjects were enrolled and 19% subjects completed the study through 9 menstrual cycles. A total of 2.1% of the subjects withdrew due to adverse events. The total exposure to 3900 mg/day tranexamic acid USP tablets in this study was 1,956 cycles. The average duration of tranexamic acid USP tablets use was 3.5 days per cycle.

The types and severity of adverse events in these two long-term open-label trials were similar to those observed in the double-blind, placebo-controlled studies although the percentage of subjects reporting them was greater in the 27-month study, most likely because of the longer study duration.

A case of severe allergic reaction to tranexamic acid USP tablets was reported in the extension trial, involving a subject on her fourth cycle of treatment, who experienced dyspnea, tightening of her throat, and facial flushing that required emergency medical treatment.

6.2 Postmarketing Experience

The following adverse reactions have been identified from postmarketing experience with tranexamic acid. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Based on US and worldwide postmarketing reports, the following have been reported in patients receiving tranexamic acid for various indications:

- Nausea, vomiting, and diarrhea
- Allergic skin reactions
- Anaphylactic shock and anaphylactoid reactions
- Thromboembolic events (e.g., deep vein thrombosis, pulmonary embolism, cerebral thrombosis, acute renal cortical necrosis, and central retinal artery and vein obstruction); cases have been associated with concomitant use of combination hormonal contraceptives
- Impaired color vision and other visual disturbances
- Dizziness

7 DRUG INTERACTIONS

No drug-drug interaction studies were conducted with tranexamic acid USP tablets.

7.1 Hormonal Contraceptives

Because tranexamic acid USP tablets are antifibrinolytic, concomitant use of hormonal contraception and tranexamic acid USP tablets may further exacerbate the increased thrombotic risk associated with combination hormonal contraceptives. For this reason, concomitant use of tranexamic acid USP tablets with combination hormonal contraceptives is contraindicated [see *Contraindications (4) and Warnings and Precautions (5.1)*].

7.2 Tissue Plasminogen Activators

Concomitant therapy with tissue plasminogen activators may decrease the efficacy of both tranexamic acid USP tablets and tissue plasminogen activators. Therefore, exercise caution if a woman taking tranexamic acid USP tablets therapy requires tissue plasminogen activators.

7.3 Factor IX Complex Concentrates or Anti-Inhibitor Coagulant Concentrates

Tranexamic acid USP Tablets are not recommended for women taking either Factor IX complex concentrates or anti-inhibitor coagulant concentrates because the risk of thrombosis may be increased [see *Warnings and Precautions (5.1) and Clinical Pharmacology (12.3)*].

7.4 All-Trans Retinoic Acid (Oral Tretinoin)

Exercise caution when prescribing tranexamic acid USP tablets to women with acute promyelocytic leukemia taking all-trans retinoic acid for remission induction because of possible exacerbation of the procoagulant effect of all-trans retinoic acid [see *Warnings and Precautions (5.1) and Clinical Pharmacology (12.3)*].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Tranexamic Acid USP Tablets are not indicated for use in pregnant women. Reproduction studies have been performed in mice, rats and rabbits and have revealed no evidence of impaired fertility or harm to the fetus due to tranexamic acid. However, tranexamic acid is known to cross the placenta and appears in cord blood at concentrations approximately equal to the maternal concentration. There are no adequate and well-controlled studies in pregnant women [see *Nonclinical Toxicology (13.1)*].

An embryo-fetal developmental toxicity study in rats and a perinatal developmental toxicity study in rats were conducted using tranexamic acid. No adverse effects were observed in either study at doses up to 4 times the recommended human oral dose of 3900 mg/day based on mg/m² (actual animal dose 1500 mg/kg/day).

8.3 Nursing Mothers

Tranexamic acid is present in the mother's milk at a concentration of about one hundredth of the corresponding serum concentration. Tranexamic Acid USP Tablets should be used during lactation only if clearly needed.

8.4 Pediatric Use

Tranexamic Acid USP Tablets are indicated for women of reproductive age and is not intended for use in premenarcheal girls.

Based on a pharmacokinetic study in 20 adolescent females, 12 to 16 years of age, no dose adjustment is needed in the adolescent population [see *Clinical Pharmacology (12.3)*].

8.5 Geriatric Use

Tranexamic Acid USP Tablets are indicated for women of reproductive age and is not intended for use by postmenopausal women.

8.6 Renal Impairment

The effect of renal impairment on the pharmacokinetics of tranexamic acid USP tablets has not been studied. Because tranexamic acid is primarily eliminated via the kidneys by glomerular filtration with more than 95% excreted as unchanged in urine, dosage adjustment in patient with renal impairment is needed [see *Dosage and Administration (2.2) and Clinical Pharmacology (12.3)*].

8.7 Hepatic Impairment


The effect of hepatic impairment on the pharmacokinetics of tranexamic acid USP tablets has not been studied. Because only a small fraction of the drug is metabolized, dosage adjustment in patients with hepatic impairment is not needed [see *Clinical Pharmacology (12.3)*].

10 OVERDOSAGE

There are no known cases of intentional overdose with tranexamic acid USP tablets and no subjects in the clinical program took more than 2 times the prescribed amount of tranexamic acid USP tablets in a 24-hour period (>7800 mg/day). However, cases of overdose of tranexamic acid have been reported. Based on these reports, symptoms of overdose may include gastrointestinal (nausea, vomiting, diarrhea); hypotensive (e.g., orthostatic symptoms); thromboembolic (arterial, venous, embolic); visual impairment; mental status changes; myoclonus; or rash. No specific information is available on the treatment of overdose with tranexamic acid USP tablets. In the event of overdose, employ the usual supportive measures (e.g., clinical monitoring and supportive therapy) as dictated by the patient's clinical status.

11 DESCRIPTION

Tranexamic Acid USP Tablets is an antifibrinolytic drug. The chemical name is trans-4-aminomethyl- clohexanecarboxylic acid. The structural formula is:



Tranexamic acid is a white crystalline powder. It is freely soluble in water and in glacial acetic acid and is very slightly soluble in ethanol and practically insoluble in ether. The molecular formula is C₈H₁₅N₂O₂ and the molecular weight is 157.2.

Tranexamic Acid USP Tablets are provided as white oval-shaped tablets and are not scored. Each tablet is debossed with the marking "FP650." The active ingredient in each tablet is 650 mg tranexamic acid.

The inactive ingredients contained in each tablet are: microcrystalline cellulose, colloidal silicon dioxide, pregelatinized corn starch, povidone, hypromellose, stearic acid, and magnesium stearate.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Tranexamic acid is a synthetic lysine amino acid derivative, which diminishes the dissolution of hemostatic fibrin by plasmin. In the presence of tranexamic acid, the lysine receptor binding sites of plasmin for fibrin are occupied, preventing binding to fibrin monomers, thus preserving and stabilizing fibrin's matrix structure.

The antifibrinolytic effects of tranexamic acid are mediated by reversible interactions at multiple binding sites within plasminogen. Native human plasminogen contains 4 to 5 lysine binding sites with low affinity for tranexamic acid (K_d = 750 μmol/L) and 1 with high affinity (K_d = 1.1 μmol/L). The high affinity lysine site of plasminogen is involved in its binding to fibrin. Saturation of the high affinity binding site with tranexamic acid displaces plasminogen from the surface of fibrin. Although plasmin may be formed by conformational changes in plasminogen, binding to and dissolution of the fibrin matrix is inhibited.

12.2 Pharmacodynamics

Tranexamic acid, at *in vitro* concentrations of 25 - 100 μM, reduces by 20 - 60% the maximal rate of plasmin lysis of fibrin catalyzed by tissue plasminogen activator (tPA).

Elevated concentrations of endometrial, uterine, and menstrual blood tPA are observed in women with heavy menstrual bleeding (HMB) compared to women with normal menstrual blood loss. The effect of tranexamic acid on lowering endometrial tPA activity and menstrual fluid fibrinolysis is observed in women with HMB receiving tranexamic acid total oral doses of 2-3 g/day for 5 days.

In healthy subjects, tranexamic acid at blood concentrations less than 10 mg/mL has no effect on the platelet count, the coagulation time or various coagulation factors in whole blood or citrated blood. Tranexamic acid, however, at blood concentrations of 1 and 10 mg/mL prolongs the thrombin time.

Cardiac Electrophysiology

The effect of tranexamic acid USP tablets on QT interval was evaluated in a randomized, single-dose, 4-way crossover study in 48 healthy females aged 18 to 49 years. Subjects received (1) tranexamic acid USP tablets 1300 mg (two 650 mg tablets), (2) tranexamic acid USP tablets 3900 mg (six 650 mg tablets; three times the recommended single dose), (3) moxifloxacin 400 mg, and (4) placebo. There was no significant increase in the corrected QT interval at any time up to 24 hours after the administration of either dose of tranexamic acid USP tablets. Moxifloxacin, the active control, was associated with a maximum 14.11 msec mean increase in corrected QT interval (moxifloxacin - placebo) at 3 hours after administration.

12.3 Pharmacokinetics

Absorption

After a single oral administration of two 650 mg tablets of tranexamic acid, the peak plasma concentration (C_{max}) occurred at approximately 3 hours (T_{max}). The absolute bioavailability of tranexamic acid USP tablets in women aged 18-49 is approximately 45%. Following multiple oral doses (two 650 mg tablets three times daily) administration of tranexamic acid USP tablets for 5 days, the mean C_{max} increased by approximately 19% and the mean area under the plasma concentration-time curve (AUC) remained unchanged, compared to a single oral dose administration (two 650 mg tablets). Plasma concentrations reached steady state at the 5th dose of tranexamic acid USP tablets on Day 2. The mean plasma pharmacokinetic parameters of tranexamic acid determined in 19 healthy women following a single (two 650 mg tablets) and multiple (two 650 mg tablets three times daily for 5 days) oral dose of tranexamic acid USP tablets are shown in Table 3.

Table 3. Mean (CV%) Pharmacokinetic Parameters Following a Single (two 650 mg tablets) and Multiple Oral Dose (two 650 mg tablets three times daily for 5 days) Administration of tranexamic acid USP tablets in 19 Healthy Women under Fasting Conditions

Parameter	Arithmetic Mean (CV%)	
	Single dose	Multiple dose
C_{max} (mcg/mL)	13.83 (32.14)	16.41 (26.19)
AUC _{0-8h} (mcg-h/mL)	77.96 (31.14)	77.67 ^a (29.39)
AUC _{inf} (mcg-h/mL)	80.19 (30.43)	-
T_{max} (h) ^b	2.5 (1 – 5)	2.5 (2 – 3.5)
$t_{1/2}$ (h)	11.08 (16.94)	-

C_{max} = maximum concentration

AUC_{0-8h} = area under the drug concentration curve from time 0 to time of last determinable concentration

AUC_{inf} = area under the drug concentration curve from time 0 to infinity T_{max} = time to maximum concentration

$t_{1/2}$ = terminal elimination half-life

^aAUC_{0-8h}(mcg-h/mL) = area under the drug concentration curve from time 0 to 8 hours

^bData presented as median (range)

Effect of food: tranexamic acid USP tablets may be administered without regard to meals. A single dose administration (two 650 mg tablets) of tranexamic acid USP tablets with food increased both C_{max} and AUC by 7% and 16%, respectively.

Distribution

Tranexamic acid is 3% bound to plasma proteins with no apparent binding to albumin. Tranexamic acid is distributed with an initial volume of distribution of 0.18 L/kg and steady-state apparent volume of distribution of 0.39 L/kg.

Tranexamic acid crosses the placenta. The concentration in cord blood after an intravenous injection of 10 mg/kg to pregnant women is about 30 mg/L, as high as in the maternal blood.

Tranexamic acid concentration in cerebrospinal fluid is about one tenth of the plasma concentration.

The drug passes into the aqueous humor of the eye achieving a concentration of approximately one tenth of plasma concentrations.

Metabolism

A small fraction of the tranexamic acid is metabolized.

Excretion

Tranexamic acid is eliminated by urinary excretion primarily via glomerular filtration with more than 95% of the dose excreted unchanged. Excretion of tranexamic acid is about 90% at 24 hours after intravenous administration of 10 mg/kg. Most elimination post intravenous administration occurred during the first 10 hours, giving an apparent elimination half-life of approximately 2 hours. The mean terminal half-life of tranexamic acid USP tablets is approximately 11 hours. Plasma clearance of tranexamic acid is 110-116 mL/min.

Specific Populations

Pregnancy (Category B)

Tranexamic Acid USP Tablets are not indicated for use in pregnant women. Tranexamic acid is known to cross the placenta and appears in cord blood at concentrations approximately equal to maternal concentration. There are no adequate and well-controlled studies in pregnant women [see *Use in Specific Populations* (8.1)].

Nursing Mothers

Tranexamic acid is present in the mother's milk at a concentration of about one hundredth of the corresponding serum concentrations. Tranexamic Acid USP Tablets should be used during lactation only if clearly needed [see *Use in Specific Populations* (8.3)].

Pediatric Use

Tranexamic Acid USP Tablets are indicated for women of reproductive age and are not intended for use in premenarcheal girls.

In a randomized, single dose, two-way crossover study of two dose levels (650 mg and 1,300 mg [two 650 mg tablets]), pharmacokinetics of tranexamic acid was evaluated in 20 female adolescents (12 to 16 years of age) with heavy menstrual bleeding. The C_{max} and AUC values after a single oral dose of 650 mg in the adolescent females were 32 – 36% less than those after a single oral dose of 1,300 mg in the adolescent females. The C_{max} and AUC values after a single oral dose of 1300 mg in the adolescent females were 20 – 25% less than those in the adult females given the same dose in a separate study. [See *Use in Specific Populations* (8.4)]

Geriatric Use

Tranexamic Acid USP Tablets are indicated for women of reproductive age and are not intended for use by postmenopausal women.

Renal Impairment

The effect of renal impairment on the disposition of tranexamic acid USP tablets has not been evaluated. Urinary excretion following a single intravenous injection of tranexamic acid declines as renal function decreases. Following a single 10 mg/kg intravenous injection of tranexamic acid in 28 patients, the 24-hour urinary fractions of tranexamic acid with serum creatinine concentrations 1.4 – 2.8, 2.8 – 5.7, and greater than 5.7 mg/dL were 51, 39, and 19%, respectively. The 24-hour tranexamic acid plasma concentrations for these patients demonstrated a direct relationship to the degree of renal impairment. Therefore, dose adjustment is needed in patients with renal impairment [see *Dosage and Administration* (2.2)].

Hepatic Impairment

The effect of hepatic impairment on the disposition of tranexamic acid USP tablets has not been evaluated. One percent and 0.5 percent of an oral dose are excreted as a dicarboxylic acid and acetylated metabolite, respectively. Because only a small fraction of the drug is metabolized, no dose adjustment is needed in patients with hepatic impairment.

Drug Interactions

No drug-drug interaction studies were conducted with tranexamic acid USP tablets.

Hormonal Contraceptives

Because tranexamic acid USP tablets are antifibrinolytic, concomitant use of hormonal contraception and tranexamic acid USP tablets may further exacerbate the increased thrombotic risk associated with combination hormonal contraceptives. For this reason, concomitant use of tranexamic acid USP tablets with combination hormonal contraceptives is contraindicated [see *Contraindications* (4), *Warnings and Precautions* (5.1) and *Drug Interactions* (7.1)]

Factor IX Complex Concentrates or Anti-inhibitor Coagulant Concentrates

Tranexamic Acid USP Tablets are not recommended in patients taking either Factor IX complex concentrates or anti-inhibitor coagulant concentrates because the risk of thrombosis may be increased [see *Warnings and Precautions* (5.1) and *Drug Interactions* (7.3)].

Tissue Plasminogen Activators

Concomitant therapy with tissue plasminogen activators may decrease the efficacy of both tranexamic acid USP tablets and tissue plasminogen activators. Therefore, exercise caution if a patient taking tranexamic acid USP tablets therapy requires tissue plasminogen activators [see *Drug Interactions* (7.2)].

All-Trans Retinoic Acid (Oral Retinoin)

In a study involving 28 patients with acute promyelocytic leukemia who were given either orally administered all-trans retinoic acid plus intravenously administered tranexamic acid, all-trans retinoic acid plus chemotherapy, or all-trans retinoic acid plus tranexamic acid plus chemotherapy, all 4 patients who were given all-trans retinoic acid plus tranexamic acid died, with 3 of the 4 deaths due to thrombotic complications. It appears that the procoagulant effect of all-trans retinoic acid may be exacerbated by concomitant use of tranexamic acid. Therefore, exercise caution when prescribing tranexamic acid USP tablets to patients with acute promyelocytic leukemia taking all-trans retinoic acid [see *Warnings and Precautions* (5.1) and *Drug Interactions* (7.4)].

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Carcinogenicity studies with tranexamic acid in male mice at doses as high as 6 times the recommended human dose of 3900 mg/day showed an increased incidence of leukemia which may have been related to treatment. Female mice were not included in this experiment.

The dose multiple referenced above is based on body surface area (mg/m²). Actual daily dose in mice was up to 5000 mg/kg/day in food.

Hyperplasia of the biliary tract and cholangioma and adenocarcinoma of the intrahepatic biliary system have been reported in one strain of rats after dietary administration of doses exceeding the maximum tolerated dose for 22 months. Hyperplastic, but not neoplastic, lesions were reported at lower doses. Subsequent long-term dietary administration studies in a different strain of rat, each with an exposure level equal to the maximum level employed in the earlier experiment, have failed to show such hyperplastic/neoplastic changes in the liver.

Mutagenesis

Tranexamic acid was neither mutagenic nor clastogenic in the *in vitro* Bacterial Reverse Mutation Assay (Ames test), *in vitro* chromosome aberration test in Chinese hamster cells, and in *in vivo* chromosome aberration tests in mice and rats.

Impairment of Fertility

Reproductive studies performed in mice, rats and rabbits have not revealed any evidence of impaired fertility or adverse effects on the fetus due to tranexamic acid.

In a rat embryo-fetal developmental toxicity study, tranexamic acid had no adverse effects on embryo-fetal development when administered during the period of organogenesis (from gestation days 6 through 17) at doses 1, 2 and 4 times the recommended human oral dose of 3900 mg/day. In a perinatal-postnatal study in rats, tranexamic acid had no adverse effects on pup viability, growth or development when administered from gestation day 6 through postnatal day 20 at doses 1, 2 and 4 times the recommended human oral dose of 3900 mg/day.

The dose multiples referenced above are based on body surface area (mg/m²). Actual daily doses in rats were 300, 750 or 1500 mg/kg/day.

13.2 Animal Toxicology and/or Pharmacology

Ocular Effects

In a 9-month toxicology study, dogs were administered tranexamic acid in food at doses of 0, 200, 600, or 1200 mg/kg/day. These doses are approximately 2, 5, and 6 times, respectively, the recommended human oral dose of 3900 mg/day based on AUC. At 6 times the human dose, some dogs developed reversible reddening and gelatinous discharge from the eyes. Ophthalmologic examination revealed reversible changes in the nictitating membrane/conjunctiva. In some female dogs, the presence of inflammatory exudate over the bulbar conjunctival mucosa was observed. Histopathological examinations did not reveal any retinal alteration. No adverse effects were observed at 5 times the human dose.

In other studies, focal areas of retinal degeneration were observed in cats, dogs and rats following oral or intravenous tranexamic acid doses at 6-40 times the recommended usual human dose based on mg/m² (actual animal doses between 250-1600 mg/kg/day).

14 CLINICAL STUDIES

The efficacy and safety of tranexamic acid USP tablets in the treatment of heavy menstrual bleeding (HMB) was demonstrated in one 3-cycle treatment and one 6-cycle treatment, randomized, double-blind, placebo-controlled study [see *Adverse Reactions* (6.1)]. In these studies, HMB was defined as an average menstrual blood loss of ≥ 80 mL as assessed by alkaline hematin analysis of collected sanitary products over two baseline menstrual cycles. Subjects were 18 to 49 years of age with a mean age of approximately 40 years, had cyclic menses every 21-35 days, and a BMI of approximately 32 kg/m². On average, subjects had an HMB history of approximately 10 years and 40% had fibroids as determined by transvaginal ultrasound. Approximately 70% were Caucasian, 25% were Black, and 5% were Asian, Native American, Pacific Islander, or Other. Seven percent (7%) of all subjects were of Hispanic origin.

In these studies, the primary outcome measure was menstrual blood loss (MBL), measured using the alkaline hematin method. The endpoint was change from baseline in MBL, calculated by subtracting the mean MBL during treatment from the mean pretreatment MBL.

The key secondary outcome measures were based on specific questions concerning limitations in social or leisure activities (LSLA) and limitations in physical activities (LPA). Large stains (soiling beyond the undergarment) were also included as a key secondary outcome measure.

14.1 Three-Cycle Treatment Study

This study compared the effects of two doses of tranexamic acid USP tablets (1950 mg and 3900 mg given daily for up to 5 days during each menstrual period) versus placebo on MBL over a 3-cycle treatment duration. Of the 294 evaluable subjects, 115 tranexamic acid USP tablets 1950 mg/day subjects, 112 tranexamic acid USP tablets 3900 mg/day subjects and 67 placebo subjects took at least one dose of study drug and had post-treatment data available.

Results are shown in Table 4. MBL was statistically significantly reduced in patients treated with 3900 mg/day tranexamic acid USP tablets compared to placebo. Study success also required achieving a reduction in MBL that was determined to be clinically meaningful to the subjects. The 1950 mg/day tranexamic acid USP tablets dose did not meet the criteria for success.

Table 4. Mean Reduction from Baseline in MBL

Treatment Arm	N	Baseline Mean MBL (mL)	Least Squares Mean Reduction in MBL (mL)	Percent Reduction in MBL
Tranexamic acid 3900 mg/day	112	169	65*	39%
Tranexamic acid 1950 mg/day	115	178	44	25%
Placebo	67	154	7	5%

* p<0.001 versus placebo

Tranexamic Acid USP Tablets also statistically significantly reduced limitations on social, leisure, and physical activities in the 3900 mg/day dose group compared to placebo (see Table 5). No statistically significant treatment difference was observed in response rates on the number of large stains.

Table 5: Secondary Outcomes in 3-Cycle Study

Outcome Measure	N	Baseline Mean ^a	Least Squares Mean Reduction ^b
Social and Leisure Activities			
3900 mg/day tranexamic acid	112	3.00	0.98 ^c
Placebo	66	2.85	0.39
Physical Activities			
3900 mg/day tranexamic acid	112	3.07	0.94 ^c
Placebo	66	2.96	0.34
Reduction in Large Stains			
3900 mg/day tranexamic acid	111		64% ^e
Placebo	67		52%

^aResponse categories: 1=not at all limited; 2=slightly limited; 3=moderately limited; 4=quite a bit limited; 5=extremely limited

^bPositive means reflect an improvement from baseline.

^cp-value <0.05 versus placebo

^dResponders are defined as subjects who experienced a reduction from baseline in frequency of large stains.

^eNon-significant difference versus placebo

14.2 Six-Cycle Treatment Study

This study compared the effects of tranexamic acid USP tablets 3900 mg/day given daily for up to 5 days during each menstrual period versus placebo on MBL over a 6-cycle treatment duration. Of the 187 evaluable subjects, 115 tranexamic acid USP tablets subjects and 72 placebo subjects took at least one dose of study drug and had post-treatment data available. Results are shown in Table 6. MBL was statistically significantly reduced in patients treated with 3900 mg/day tranexamic acid USP tablets compared to placebo. Study success also required achieving a reduction in MBL that was determined to be clinically meaningful to the subjects.

Table 6. Mean Reduction from Baseline in MBL

Treatment Arm	N	Baseline Mean MBL (mL)	Least Squares Mean Reduction in MBL (mL)	Percent Reduction in MBL
Tranexamic acid 3900 mg/day	115	172	66*	38%
Placebo	72	153	18	12%

* p<0.001 versus placebo

Limitations on social, leisure, and physical activities were also statistically significantly reduced in the tranexamic acid USP tablets group compared to placebo (see Table 7). No statistically significant treatment difference was observed in response rates on the number of large stains.

Table 7: Secondary Outcomes in 6-Cycle Study

Outcome Measure	N	Baseline Mean ^a	Least Squares Mean Reduction ^b
Social and Leisure Activities			
3900 mg/day tranexamic acid	115	2.92	0.85 ^c
Placebo	72	2.74	0.44
Physical Activities			
3900 mg/day tranexamic acid	115	3.05	0.87 ^c
Placebo	72	2.90	0.40
		N	Responders^d
Reduction in Large Stains			
3900 mg/day tranexamic acid	115		57% ^e
Placebo	72		51%

^aResponse categories: 1=not at all limited; 2=slightly limited; 3=moderately limited; 4=quite a bit limited; 5=extremely limited

^bPositive means reflect an improvement from baseline.

^cp-value <0.05 versus placebo


^dResponders are defined as subjects who experienced a reduction from baseline in frequency of large stains.

^eNon-significant difference versus placebo

14.3 MBL Results over Time

The efficacy of tranexamic acid USP tablets 3900 mg/day over 3 menstrual cycles and over 6 menstrual cycles was demonstrated versus placebo in the double-blind, placebo-controlled efficacy studies (see Figure 1). The change in MBL from baseline was similar across all post-baseline treatment cycles.

Figure 1: MBL Levels over Duration of Therapy



16 HOW SUPPLIED/STORAGE AND HANDLING

Tranexamic Acid USP Tablets are provided as white oval-shaped tablets. Each tablet is debossed with the marking "FP650" and are supplied as:

Quantity	Package Type	NDC Number
30 tablets	HDPE bottle	69918-301-30

Storage

Store at room temperature 25° C (77° F); excursions permitted to 15-30° C (59-86° F). [See USP controlled Room Temperature]

17 PATIENT COUNSELING INFORMATION

See FDA-approved patient labeling (Patient Information)

Instruct patients that the usual schedule is to take two tablets with liquids, three times a day during menstruation. Patients should be instructed not to exceed 3 doses (6 tablets) in a 24-hour period or to take for more than 5 days in any menstrual cycle.

Inform patients that they should immediately stop tranexamic acid USP tablets if they notice any eye symptoms or change in their vision. Instruct them to report any such problems promptly to their physician and to follow-up with an ophthalmologist for a complete ophthalmic evaluation, including dilated retinal examination of the retina.

Inform patients that they should stop tranexamic acid USP tablets and seek immediate medical attention if they notice symptoms of a severe allergic reaction (e.g., shortness of breath or throat tightening).

Instruct patients that common side effects of tranexamic acid USP tablets include headache, sinus and nasal symptoms, back pain, abdominal pain, musculoskeletal pain, joint pain, muscle cramps, migraine, anemia and fatigue.

Advise patients to contact their healthcare provider if their heavy menstrual bleeding symptoms persist or worsen.

Remind patients to read the Patient Labeling carefully.

PATIENT INFORMATION

TRANEXAMIC ACID, USP TABLETS

Read the Patient Information that comes with tranexamic acid USP tablets before you start using the drug and each time you get a refill. There may be new information. This leaflet does not take the place of talking with your healthcare provider about your medical condition or your treatment.

What is tranexamic acid USP tablets?

Tranexamic Acid USP Tablets is a prescription medicine used to treat your heavy monthly period (menstruation) when your bleeding gets in the way of social, leisure and physical activities. Tranexamic Acid USP Tablets does not contain any hormones. On average, tranexamic acid USP tablets has been shown to lower the amount of blood lost during your monthly period by about one-third, but it is not meant to stop your period.

Tranexamic Acid USP Tablets is taken only during your period and is not meant to treat pre-menstrual symptoms (symptoms that occur before your bleeding starts). Tranexamic Acid USP Tablets does not affect your fertility and cannot be used as birth control. Tranexamic Acid USP Tablets does not protect you against diseases that you may get if you have unprotected sex.

Tranexamic Acid USP Tablets has not been studied in adolescents younger than 18 years of age.

Tranexamic Acid is not for women who have already gone through menopause (post-menopausal).

Who should not take tranexamic acid USP tablets?

Do not take tranexamic acid USP tablets if you:

- Are using a form of birth control that contains estrogen and a progestin (like a birth control pill, patch, or vaginal ring). Ask your healthcare provider before taking tranexamic acid USP tablets if you are not sure if your birth control method contains estrogen and a progestin.
- Currently have a blood clot
- Have ever had a blood clot
- Have been told that you are at risk of having a blood clot
- Are allergic to tranexamic acid

What should I tell my healthcare provider before taking tranexamic acid USP tablets?

Before taking tranexamic acid USP tablets, tell your healthcare provider about all of your medical conditions, including whether:

- You have ever had a blood clot or been told that you are at risk of having a blood clot
- You are using a form of birth control that contains estrogen and a progestin (like a birth control pill, patch, or vaginal ring). Using hormonal birth control along with tranexamic acid USP tablets may increase your chance of having a serious blood clot, stroke, or heart attack. For this reason, do not use tranexamic acid USP tablets if you use a form of birth control that contains estrogen and a progestin.
- You are pregnant or think you may be pregnant
- You are breastfeeding or plan to breast-feed. Tranexamic Acid USP Tablets can pass into your milk. Talk to your healthcare provider about the best way to feed your baby if you take tranexamic acid USP tablets.
- The time between the start of your periods is less than 21 days or more than 35 days
- You have any other medical conditions

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Tranexamic Acid USP Tablets and other medicines can affect each other, causing side effects. Tranexamic Acid USP Tablets can affect the way other medicines work and other medicines can affect how tranexamic acid USP tablets work.

Especially tell your healthcare provider if you take:

- Birth control pills or other hormonal birth control
- Medicines used to help your blood form clots
- Medicines used to break up blood clots
- Any medicines to treat leukemia

Ask your healthcare provider if you are not sure if your medicine is one that is described above.

How should I take tranexamic acid USP tablets?

- Take tranexamic acid USP tablets exactly as your healthcare provider tells you.
- Do not take tranexamic acid USP tablets until your period has started.
- Do not take tranexamic acid USP tablets for more than 5 days in a row.
- Do not take tranexamic acid USP tablets when you do not have your period.
- Once your period has started, take 2 tablets of tranexamic acid USP tablets three times per day (e.g., in the morning, afternoon, and evening).
- Tranexamic Acid USP Tablets should be swallowed whole and not chewed or broken apart.
- Tranexamic Acid USP Tablets may be taken with or without food.
- Do not take more than 6 tablets of tranexamic acid USP tablets in a day. If you take more than 6 tablets, call your healthcare provider.
- If you miss a dose, take it when you remember, and then take your next dose at least six hours later. Do not take more than two tablets at a time to make up for missed doses.
- If tranexamic acid USP tablets does not help to lessen bleeding with your periods after 2 cycles or seems to stop working, talk to your healthcare provider.

What are the possible side effects of tranexamic acid USP tablets? Tranexamic Acid USP Tablets can cause serious side effects, including:

- Blood clots. You may have a higher risk of having serious blood clots if you take tranexamic acid USP tablets with:
 - medicines used to help your blood form clots
 - some medicines used to treat leukemia
- Eye changes. Stop taking tranexamic acid USP tablets and promptly report any eye problems you have while taking tranexamic acid USP tablets. Your doctor will refer you to an eye doctor who will examine your eyes.
- Allergic reaction. If you have severe shortness of breath and your throat feels tight, stop taking tranexamic acid USP tablets and get medical care right away.

The most common side effects of tranexamic acid USP tablets include:

- Headaches
- Sinus and nasal problems
- Back pain
- Pain in your abdomen
- Pain in your muscles or joints
- Anemia
- Fatigue

Tell your healthcare provider if you have any side effect that bothers you or does not go away.

These are not all of the possible side effects of tranexamic acid USP tablets. For more information, ask your healthcare provider or pharmacist.

If you notice a change in your usual bleeding pattern that worries you, or your heavy bleeding continues, contact your healthcare provider right away. This may be a sign of a more serious condition.

Call your healthcare provider for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. You may also report side effects to Amring Pharmaceuticals Inc. at 1-844-Amring1 (1-844-267-4641).

How should I store tranexamic acid USP tablets?

Store tranexamic acid USP tablets at room temperature between 59°F to 86°F (15°C to 30°C).

Keep tranexamic acid USP tablets and all medicines out of the reach of children.

General information about tranexamic acid USP tablets

Medicines are sometimes prescribed for conditions that are not mentioned in Patient Information Leaflets. Do not use tranexamic acid USP tablets for a condition for which it was not prescribed. Do not give tranexamic acid USP tablets to other people, even if they have the same symptoms that you have. It may harm them.

This patient information leaflet summarizes the most important information about tranexamic acid USP tablets. If you would like more information about tranexamic acid USP tablets, talk with your healthcare provider. You can ask your healthcare provider or pharmacist for information about tranexamic acid USP tablets that is written for healthcare professionals. For more information, call 1-844-Amring1 (1-844-267-4641).

What are the ingredients of tranexamic acid USP tablets?

Active ingredient: tranexamic acid

Inactive ingredients: microcrystalline cellulose, colloidal silicon dioxide, pregelatinized corn starch, povidone, hypromellose, stearic acid, and magnesium stearate.

This Patient Information has been approved by the U.S. Food and Drug Administration.

Manufactured for:
Amring Pharmaceuticals Inc.
Berwyn, PA 19312



Manufactured by:
Mikart, LLC.
Atlanta, GA 30318

Rev. 03/2019

1221E00